

Package: geodata (via r-universe)

September 14, 2026

Type Package

Title Access Geographic Data

Version 0.6-10

Date 2026-04-05

Depends R (>= 3.5.0), terra (>= 1.6.41)

Imports rappdirs

Suggests jsonlite, R.utils, httr, archive

Encoding UTF-8

Maintainer Robert J. Hijmans <r.hijmans@gmail.com>

Description Functions for downloading of geographic data for use in spatial analysis and mapping. The package facilitates access to climate, crops, elevation, land use, soil, species occurrence, accessibility, administrative boundaries and other data.

License GPL (>=3)

URL <https://rspatial.github.io/geodata/>

BugReports <https://github.com/rspatial/geodata/issues/>

Config/pak/sysreqs libgdal-dev gdal-bin libgeos-dev libproj-dev libsqlite3-dev

Repository <https://predictiveecology.r-universe.dev>

Date/Publication 2026-06-16 21:48:07 UTC

RemoteUrl <https://github.com/rspatial/geodata>

RemoteRef HEAD

RemoteSha fe495b93cd1fb54d737f1fab5ce1ed160660458b

Contents

geodata-package	2
bio_oracle	3
cmip6	4
country_codes	5

crop_calendar_sacks	6
crop_monfreda	7
crop_spam	8
cropland	9
elevation	10
footprint	11
gadm	12
geodata_path	13
landcover	13
osm	14
population	15
rice_calendar	16
soil_af	16
soil_af_elements	18
soil_af_isda	19
soil_af_isda_vsi	21
soil_af_water	22
soil_world	24
soil_world_vsi	25
sp_occurrence	27
travel_time	29
world	30
worldclim	31

Index	33
--------------	-----------

geodata-package	<i>Access Geographic Data</i>
-----------------	-------------------------------

Description

Functions for downloading of geographic data for use in spatial analysis and mapping. The package facilitates access to climate, crops, elevation, land use, soil, species occurrence, accessibility, administrative boundaries and other data.

Function	Description
bio_oracle	Marine data from bio-oracle
cmip6_world	Downscaled and calibrated CMIP6 projected future climate data
cmip6_tile	Downscaled and calibrated CMIP6 data by tile
country_codes	Country codes
crop_calendar_sacks	Sachs crop calendar data
crop_monfreda	Monfreda crop data (area, yield)
crop_spam	SPAM crop data (area, yield, value)
cropland	Cropland density for the world from three sources
elevation_3s	Elevation data for tile (3 seconds resolution)
elevation_30s	Elevation data for by country (30 seconds resolution)
elevation_global	Global elevation data (various resolutions)
gadm	Administrative boundaries for any country in the world

world	Boundaries for the countries in the world
landcover	Global landcover data
footprint	Human footprint data
osm	OpenStreetMap data by country
population	Download population density data
soil_af	Chemical and physical soil properties data for Africa for different soil depths
soil_af_water	Physical soil properties for Africa for water balance computations
soil_af_elements	Soil element concentration data for Africa
soil_af_isda	Soil data for Africa derived from the iDSA data set
soil_world_vsi	Virtually connect to the global soilgrids data
soil_world	Global soils data
sp_occurrence	Species occurrence data from the Global Biodiversity Information Facility
travel_time	Travel time to cities and ports
worldclim_global	Global climate data
worldclim_country	Climate data by country
worldclim_tile	Climate data by tile

 bio_oracle

Marine data

Description

Marine data from Bio-Oracle

Usage

```
bio_oracle(path, var, stat, benthic=FALSE,
           depth="Mean", time="Present", rcp, ...)
```

Arguments

<code>path</code>	character. Path for storing the downloaded data. See geodata_path
<code>var</code>	character. Variable of interest. One of 'Calcite', 'Chlorophyll', 'Cloud.cover', 'Current.Velocity', 'Diffuse.attenuation', 'Dissolved.oxygen', 'Ice.cover', 'Ice.thickness', 'Iron', 'Light.bottom', 'Nitrate', 'Par', 'pH', 'Phosphate', 'Phytoplankton', 'Primary.productivity', 'Salinity', 'Silicate', 'Temperature'
<code>stat</code>	character. Statistic of interest. One of 'Lt.max', 'Lt.min', 'Max', 'Mean', 'Min', 'Range'. It should be "" if var is "pH"
<code>benthic</code>	logical. If FALSE surface data are returned
<code>depth</code>	character. Either "Min", "Mean", or "Max". Only relevant if benthic is TRUE
<code>time</code>	character. Either "Present", "2050" or "2100"
<code>rcp</code>	character. Either "26", "45", "60", or "85"
<code>...</code>	additional arguments passed to download.file

Value

SpatRaster

References

Assis, J., Tyberghein, L., Bosh, S., Verbruggen, H., Serrão, E.A., & De Clerck, O. (2017). Bio-ORACLE v2.0: Extending marine data layers for bioclimatic modelling. *Global Ecology and Biogeography* 27: 277-284.

See Also

<https://bio-oracle.org/>

Examples

```
# this is a large download
x <- bio_oracle(path=tempdir(), "Salinity", "Max",
  benthic=TRUE, depth="Mean", time="Present")
```

cmip6	<i>CMIP6 climate model data</i>
-------	---------------------------------

Description

Download downscaled and calibrated CMIP6 climate data for projected future climates. Either for the entire world or for a 30 degrees tile. For more information see <https://www.worldclim.org/>

Usage

```
cmip6_world(model, ssp, time, var, res, path, ...)

cmip6_tile(lon, lat, model, ssp, time, var, path, ...)
```

Arguments

model	character. Climate model abbreviation. One of "ACCESS-CM2", "ACCESS-ESM1-5", "AWI-CM-1-1-MR", "BCC-CSM2-MR", "CanESM5", "CanESM5-CanOE", "CMCC-ESM2", "CNRM-CM6-1", "CNRM-CM6-1-HR", "CNRM-ESM2-1", "EC-Earth3-Veg", "EC-Earth3-Veg-LR", "FIO-ESM-2-0", "GFDL-ESM4", "GISS-E2-1-G", "GISS-E2-1-H", "HadGEM3-GC31-LL", "INM-CM4-8", "INM-CM5-0", "IPSL-CM6A-LR", "MIROC-ES2L", "MIROC6", "MPI-ESM1-2-HR", "MPI-ESM1-2-LR", "MRI-ESM2-0", "UKESM1-0-LL"
ssp	character. A valid Shared Socio-economic Pathway code: "126", "245", "370" or "585"
time	character. A valid time period. One of "2021-2040", "2041-2060", or "2061-2080"

var	character. Variables name. One of "tmin", "tmax", "prec" and "bioc"
res	numeric. Spatial resolution. One of 10, 5, 2.5 and 0.5 (minutes of a degree). Silently ignored if provided to <code>cmip6_tile()</code> , available only at <code>res=0.5</code>
path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file
lon	numeric. Longitude
lat	numeric. Latitude

Value

SpatRaster

See Also[vrt](#) to combine tiles**Examples**

```
# download of large files takes a while
tmin10 <- cmip6_world("CNRM-CM6-1", "585", "2061-2080",
  var="tmin", res=10, path=tempdir())
```

country_codes	<i>Get country codes</i>
---------------	--------------------------

Description

Get country codes for all countries in the world.

Usage

```
country_codes(query=NULL)
```

Arguments

query character. A single word that can be used to subset the returned data.frame

Value

data.frame

Examples

```
cc <- country_codes()
head(cc)

p <- country_codes(query="Per")
p
```

crop_calendar_sacks	<i>Sacks crop calendar data</i>
---------------------	---------------------------------

Description

Download Sacks crop calendar data. The crops available are returned by `sacksCrops`

Usage

```
crop_calendar_sacks(crop="", path, ...)
```

```
sacksCrops()
```

Arguments

<code>crop</code>	character. Crop name. See <code>sacksCrops</code> for valid names
<code>path</code>	character. Path for storing the downloaded data. See geodata_path
<code>...</code>	additional arguments passed to download.file

Value

`SpatRaster`

References

Sacks, W.J., D. Deryng, J.A. Foley, and N. Ramankutty, 2010. Crop planting dates: an analysis of global patterns. *Global Ecology and Biogeography* 19: 607-620. doi:10.1111/j.1466-8238.2010.00551.x.

See Also

<https://sage.nelson.wisc.edu/data-and-models/datasets/crop-calendar-dataset/>

Examples

```
# download may take > 5s
cas <- crop_calendar_sacks("cassava", path=tempdir())
```

crop_monfreda	<i>Monfreda crop data</i>
---------------	---------------------------

Description

Monfreda global crop data (area, yield) for 175 crops.

Data may be freely used for research, study, or teaching, but must be cited appropriately (see below).

Re-release of the data, or incorporation of the data into a commercial product, is allowed only with explicit permission.

Usage

```
monfredaCrops()
crop_monfreda(crop="", var="area_ha", path, ...)
```

Arguments

crop	character. Crop name(s). See <code>monfredaCrops</code> for valid names
var	character. The variable(s) of interest. Choose from "area_ha" (crop area in ha per cell), "area_f" (crop area as a fraction of each cell), "area_q" (quality of the crop area data), "yield" (crop yield in Mg/ha), "yield_q" (quality of the yield data), "prod" (production per grid cell in Mg), or "all"
path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file

Value

SpatRaster

References

Monfreda, C., N. Ramankutty, and J. A. Foley (2008), Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, *Global Biogeochem. Cycles*, 22, GB1022, doi:10.1029/2007GB002947.

See Also

<http://www.earthstat.org/harvested-area-yield-175-crops/>

Examples

```
# download may take > 5s
mcas <- crop_monfreda("cassava", path=tempdir())
mcas
names(mcas)
```

crop_spam

SPAM crop data

Description

SPAM crop data. For each of 42 crops or crop groups get a 10-minute spatial resolution raster with the crop area, yield, production or value by cropping system (rainfed or irrigated, and subsistence, low-input or high-input).

The global data are for 2010. The Africa dataset is for 2017.

Usage

```
spamCrops()
crop_spam(crop="", var="area", path, africa=FALSE, ...)
```

Arguments

crop	character. See spamCrops for valid names
var	character. variable of interest. Must be one of "yield", "harv_area" (harvested area), "phys_area" (physical area), "prod" (production) or "val_prod" (value of production)
path	character. Path for storing the downloaded data. See geodata_path
africa	logical. retrieve the (more up to date) data for Africa instead of global data
...	additional arguments passed to download.file

Value

SpatRaster

References

International Food Policy Research Institute, 2019. Global Spatially-Disaggregated Crop Production Statistics Data for 2010 Version 2.0. <https://doi.org/10.7910/DVN/PRFF8V>, Harvard Dataverse, V4.

International Food Policy Research Institute, 2020. Spatially-Disaggregated Crop Production Statistics Data in Africa South of the Sahara for 2017. <https://doi.org/10.7910/DVN/FSSKBW>, Harvard Dataverse, V3.

See Also

<https://www.mapspam.info/data/>

Examples

```
# downloads a large file
cas <- crop_spam("cassava", "area", path=tempdir(), TRUE)
```

cropland	<i>Cropland distribution data</i>
----------	-----------------------------------

Description

Cropland distribution data at a 30-seconds spatial resolution from three sources:

worldcover is derived from the ESA WorldCover data set at 0.3-seconds resolution. (License CC BY 4.0), see <https://esa-worldcover.org/en>. Values were aggregated and represent the fraction cropland in each cell.

glad is derived from the "Global cropland expansion in the 21st century" (Potatov et al) data available [here](#). Values were aggregated and resampled. They represent the fraction cropland in each cell. There are five layers representing the following years: 2003, 2007, 2011, 2015, and 2019.

QED has cropland distribution data for Africa. The values are probabilities of cropland presence estimated with a neural network that was trained on an initial 1-million point survey conducted in 2015. License: CC-BY-SA 4.0; <https://about.maps.qed.ai/>

Usage

cropland(source, path, year, ...)

Arguments

- source character. One of "WorldCover", "GLAD", or "QED"
- path character. Path for storing the downloaded data. See [geodata_path](#)
- year numeric. Optional for the GLAD dataset to get data for a single year. One of 2003, 2007, 2011, 2015, and 2019
- ... additional arguments passed to [download.file](#)

Value

SpatRaster

References

WorldCover: Zanaga, D., Van De Kerchove, R., De Keersmaecker, W., Souverijns, N., Brockmann, C., Quast, R., Wevers, J., Grosu, A., Paccini, A., Vergnaud, S., Cartus, O., Santoro, M., Fritz, S., Georgieva, I., Lesiv, M., Carter, S., Herold, M., Li, Linlin, Tsendbazar, N.E., Ramoino, F., Arino, O., 2021. ESA WorldCover 10 m 2020 v100. doi:10.5281/zenodo.5571936.

GLAD: Potapov, P., S. Turubanova, M.C. Hansen, A. Tyukavina, V. Zalles, A. Khan, X.-P. Song, A. Pickens, Q. Shen, J. Cortez, 2021. Global maps of cropland extent and change show accelerated cropland expansion in the twenty-first century. Nature Food. doi:10.1038/s43016-021-00429-z

See Also

[landcover](#)

elevation

Elevation

Description

Elevation data for any country. The main data source is Shuttle Radar Topography Mission (SRTM) , specifically the hole-filled CGIAR-SRTM (90 m resolution) from <https://srtm.csi.cgiar.org/>. These data are only available for latitudes between -60 and 60.

The 1 km (30 arc seconds) data were aggregated from SRTM 90 m resolution data and supplemented with the GTOPO30 data for high latitudes (>60 degrees).

Usage

```
elevation_3s(lon, lat, path, ...)
elevation_30s(country, path, mask=TRUE, subs="", ...)
elevation_global(res, path, ...)
```

Arguments

lon	numeric. Longitude
lat	numeric. Latitude
path	character. Path for storing the downloaded data. See geodata_path
country	character. Country name or code
mask	logical. set grid cells outside of the country boundaries to NA
subs	character
res	numeric. Valid resolutions are 10, 5, 2.5, and 0.5 (minutes of a degree)
...	additional arguments passed to download.file

Value

SpatRaster

Examples

```
be <- elevation_30s(country="BEL", path=tempdir() )
```

footprint

Human footprint

Description

The "human footprint" is an estimate of the direct and indirect human pressures on the environment. The human pressure is measured using eight variables including built-up environments, population density, electric power infrastructure, crop lands, pasture lands, roads, railways, and navigable waterways. It is expressed on a scale of 0 (low) to 50 (high footprint).

See <https://www.nature.com/articles/sdata201667> for the details.

The original data are available here:

<https://sedac.ciesin.columbia.edu/data/collection/wildareas-v3>

Data are available for two years: 1993 and 2009, for all terrestrial areas except Antarctica. The footprint of seas and oceans was set to zero. The original data was in the Mollweide projection at a 1000 m spatial resolution. The data available through this function was transformed to a longitude/latitude grid at 30-seconds resolution.

Users are free to use, copy, distribute, transmit, and adapt the work for commercial and non-commercial purposes, without restriction, as long as clear attribution of the source is provided.

Usage

```
footprint(year=2009, path, ...)
```

Arguments

year	character. "1993" or "2009"
path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file

Value

SpatRaster

References

Venter, O., E. W. Sanderson, A. Magrath, J. R. Allan, J. Beher, K. R. Jones, H. P. Possingham, W. F. Laurance, P. Wood, B. M. Fekete, M. A. Levy, and J. E. Watson. 2016. Sixteen Years of Change in the Global Terrestrial Human Footprint and Implications for Biodiversity Conservation. *Nature Communications* 7:12558. <https://doi.org/10.1038/ncomms12558>.

See Also

[landcover](#)

gadm*Administrative boundaries*

Description

Get administrative boundaries for any country in the world. Data are read from files that are downloaded if necessary.

Usage

```
gadm(country, level=1, path, version="latest", resolution=1, ...)
```

Arguments

country	character. Three-letter ISO code or full country name. If you provide multiple names they are all downloaded and rbind-ed together
level	numeric. The level of administrative subdivision requested. (starting with 0 for country, then 1 for the first level of subdivision)
path	character. Path for storing the downloaded data. See geodata_path
version	character. Either "latest" or GADM version number (can be "3.6", "4.0" or "4.1")
resolution	integer indicating the level of detail. Only for version 4.1. It should be either 1 (high) or 2 (low)
...	additional arguments passed to download.file

Details

The data are from <https://gadm.org>

Value

SpatVector

See Also

[world](#)

Examples

```
bel <- gadm(country="BEL", level=1, path=tempdir())
```

geodata_path	<i>Set the data path</i>
--------------	--------------------------

Description

This function allows you set or get the default download cache path for the geodata package. The default path is ignored if you set the path argument in a geodata function to another value.

A special value is "user_data_dir". This sets it to the location returned by `file.path(rappdirs::user_data_dir(), "geodata")`

Alternatively, you can also set a system variable "GEODATA_PATH" to the desired path.

Usage

```
geodata_path(path, persistent=TRUE)
```

Arguments

path	character. Path name where geodata should be download and cache data. If missing, the current default path is returned. Use NA to restore the original value
persistent	logical. Should the path be persistent (automatically used in future R sessions)?

Value

character

Examples

```
geodata_path()
```

landcover	<i>Landcover data</i>
-----------	-----------------------

Description

Landcover data at 30-seconds spatial resolution for (most of) the world. Values are the fraction of a landcover class in each cell. The values are derived from the ESA WorldCover data set at 0.3-seconds resolution. (License CC BY 4.0). See <https://esa-worldcover.org/en> for more information.

Usage

```
landcover(var, path, ...)
```

Arguments

var	character. One of "trees", "grassland", "shrubs", "cropland", "built", "bare", "snow", "water", "wetland", "mangroves", "moss"
path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file

Value

SpatRaster

References

Zanaga, D., Van De Kerchove, R., De Keersmaecker, W., Souverijns, N., Brockmann, C., Quast, R., Wevers, J., Grosu, A., Paccini, A., Vergnaud, S., Cartus, O., Santoro, M., Fritz, S., Georgieva, I., Lesiv, M., Carter, S., Herold, M., Li, Linlin, Tsendbazar, N.E., Ramoino, F., Arino, O., 2021. ESA WorldCover 10 m 2020 v100. doi:10.5281/zenodo.5571936.

See Also

[landcover](#)

osm

OpenStreetMap data

Description

Get OpenStreetMap (OSM) data

Usage

```
osm(country, var, path, proxy=FALSE, ...)
```

Arguments

country	character. Three-letter ISO code or full country name
var	character. Currently it can be one of "places", "highways", or "railway"
path	character. Path for storing the downloaded data. See geodata_path
proxy	logical. Return a SpatVectorProxy?
...	additional arguments passed to download.file

Details

License: Open Data Commons Open Database License (ODbL).

See <https://www.openstreetmap.org/copyright>

Value

SpatVector

Examples

```
aruba <- osm(country="Aruba", "places", path=tempdir())
```

 population

population density

Description

Download population density data.

Source: Gridded Population of the World (GPW), v4. Documentation:

<http://sedac.ciesin.columbia.edu/data/collection/gpw-v4/documentation>

Usage

```
population(year, res, path, ...)
```

Arguments

year	numeric. One of 2000, 2005, 2010, 2015, 2020
res	numeric. Valid resolutions are 10, 5, 2.5, and 0.5 (minutes of a degree)
path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file

Value

SpatRaster

References

Center for International Earth Science Information Network - CIESIN - Columbia University. 2018. Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 11. Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC). doi:10.7927/H49C6VHW. Accessed 6 July 2021.

Examples

```
# download may take > 5s
pop <- population(2020, 10, path=tempdir())
```

rice_calendar	<i>crop calendar for rice</i>
---------------	-------------------------------

Description

Get crop calendar and production data for rice

Usage

```
rice_calendar(path, ...)
```

Arguments

path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file

Value

SpatVectorCollection

References

Laborte, A.G.; Gutierrez, M.A.; Balanza, J.G.; Saito, K.; Zwart, S.J.; Boschetti, M.; Murty, MVR; Villano, L.; Aunario, J.K.; Reinke, R.; Koo, J.; Hijmans, R.J.; Nelson, A., 2017. RiceAtlas, a spatial database of global rice calendars and production. Scientific Data 4: 170074 [doi:10.1038/sdata.2017.74](#)

Examples

```
# first time api call takes a while
rice <- rice_calendar(path=tempdir())
cal <- rice[1]
```

soil_af	<i>Soil data for Africa</i>
---------	-----------------------------

Description

Download chemical soil properties data for Africa for different soil depths. The spatial resolution is 30 arc-seconds (about 1 km²), aggregated from the original 250m resolution.

There are more recent estimations for some of the properties available in other data sets. See [soil_af_isda](#) and [soil_world](#).

For more info, see <https://web.archive.org/web/20250421195225/https://www.isric.org/projects/soil-property-maps-africa-250-m-resolution>

The data have a CC-BY 4.0 NC license

Usage

```
soil_af(var, depth, path, ...)
```

Arguments

var character. Variables name (see Details below). One of: "acid-exch", "Al-extr", "Al-exch", "bases-exch", "BDR", "BLKD", "Ca-exch", "CEC", "clay", "coarse", "drain", "ECN", "K-exch", "Mg-exch", "Na-exch", "Ntot", "pH", "sand", "silt", "SOC"

depth numeric. One of 5, 15, 30, 60, 100, 200. This is shorthand for the following depth ranges: 0-5, 5-15, 15-30, 30-60, 60-100, 100-200 cm. Or one of 20, 50 for 0-20 or 20-50 cm

path character. Path for storing the downloaded data. See [geodata_path](#)

... additional arguments passed to [download.file](#)

Details

var	description	unit
clay	Soil texture fraction clay	%
sand	Soil texture fraction sand	%
silt	Soil texture fraction silt	%
coarse	Coarse fragments volumetric	%
SOC	Organic carbon	$g \cdot kg^{-1}$ (‰)
BLKD	Bulk density (fine earth)	$kg \cdot m^{-3}$
BDR	Depth to bedrock	cm
.	.	.
pH	pH (H_2O)	-
ECN	Electrical conductivity	mS/m (?)
acid-exch	Exchangeable acidity	$cmol(+) \cdot kg^{-1}$
bases-exch	Sum of exchangeable bases	$cmol(+) \cdot kg^{-1}$
CEC	Cation Exchange Capacity	$cmol(+) \cdot kg^{-1}$
Al-extr	Extractable Aluminum (Mehlich 3)	$mg \cdot kg^{-1}$ (ppm)
Al-exch	Exchangeable Aluminum	$cmol(+) \cdot kg^{-1}$
Ca-exch	Exchangeable Calcium	$cmol(+) \cdot kg^{-1}$
K-exch	Exchangeable Potassium	$cmol(+) \cdot kg^{-1}$
Mg-exch	Exchangeable Magnesium	$cmol(+) \cdot kg^{-1}$
Na-exch	Exchangeable Sodium	$cmol(+) \cdot kg^{-1}$
Ntot	Total nitrogen	$g \cdot kg^{-1}$

Value

SpatRaster

References

Hengl T, Heuvelink GBM, Kempen B, Leenaars JGB, Walsh MG, Shepherd KD, et al. (2015) Mapping Soil Properties of Africa at 250 m Resolution: Random Forests Significantly Improve Current Predictions. PLoS ONE 10(6): e0125814. doi:10.1371/journal.pone.0125814

See Also

[soil_af_elements](#), [soil_af_isda](#), [soil_world_vsi](#)

Examples

```
# downloads a large file
aph <- soil_af(var="ph", depth=5, path=tempdir())
```

soil_af_elements	<i>Soil elements data for Africa</i>
------------------	--------------------------------------

Description

Connect to or download chemical soil element concentration (for the 0-30 cm topsoil) data for Africa. The spatial resolution is 30 arc-seconds (about 1 km²), aggregated from the original 250 m spatial resolution.

The data have an Open Data Commons Open Database License (ODbL)

For more information, see <https://web.archive.org/web/20250421195225/https://www.isric.org/projects/soil-property-maps-africa-250-m-resolution>

Usage

```
soil_af_elements(var, path, ...)
```

Arguments

var	character. Variables name. One of: "Al", "B", "Ca", "Cu", "Fe", "K", "Mg", "Mn", "N", "Na", "P", "Ptot", "Zn". See Details
path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file

Details

var	description	unit
Al	Extractable aluminum	$mg \cdot kg^{-1}$ (ppm)
B	Extractable boron	$mg \cdot kg^{-1}$ (ppm)
Ca	Extractable calcium	$mg \cdot kg^{-1}$ (ppm)
Cu	Extractable copper	$mg \cdot kg^{-1}$ (ppm)

Fe	Extractable iron	$mg \cdot kg^{-1}$ (ppm)
K	Extractable potassium	$mg \cdot kg^{-1}$ (ppm)
Mg	Extractable magnesium	$mg \cdot kg^{-1}$ (ppm)
Mn	Extractable manganese	$mg \cdot kg^{-1}$ (ppm)
N	Organic nitrogen	$mg \cdot kg^{-1}$ (ppm)
Na	Extractable sodium	$mg \cdot kg^{-1}$ (ppm)
P	Extractable phosphorus	$mg \cdot (100 \cdot kg^{-1})$
Ptot	Total phosphorus	$mg \cdot (100 \cdot kg^{-1})$
Zn	Extractable zinc	$mg \cdot kg^{-1}$ (ppm)

Value

SpatRaster

References

Hengl T, Heuvelink GBM, Kempen B, Leenaars JGB, Walsh MG, Shepherd KD, et al. (2015) Mapping Soil Properties of Africa at 250 m Resolution: Random Forests Significantly Improve Current Predictions. PLoS ONE 10(6): e0125814. doi:10.1371/journal.pone.0125814

See Also

[soil_af](#), [soil_af_isda](#), [soil_world](#)

Examples

```
# downloads a large file
fe <- soil_af_elements("Fe", path=tempdir(), quiet=TRUE)
```

soil_af_isda	<i>iSDA soil data for Africa</i>
--------------	----------------------------------

Description

Download soil data for Africa derived from the iDSA data set. The original data were aligned and aggregated to 30 arc-seconds (about 1 km²). The original spatial resolution was 30m.

For more info see:

<https://envirometrix.nl/isdasoil-open-soil-data-for-africa/>

<https://zenodo.org/search?page=1&size=20&q=iSDAsoil>

Usage

```
soil_af_isda(var, depth=20, error=FALSE, path, virtual=FALSE, ...)
```

Arguments

var	character. Name of the soil variable. Should be one of: "Al", "bdr", "clay", "C.tot", "Ca", "db.od", "eCEC.f", "Fe", "K", "Mg", "N.tot", "oc", "P", "pH.H2O", "sand", "silt", "S", "texture", "wpg2", "Zn". See the Details section below
depth	numeric. One of 20 (for 0-20 cm) and 50 (for 20-50 cm). Ignored if var="bdr" for which the depth is always 0-200 cm
error	logical. If TRUE the error estimates are returned
path	character. Path for storing the downloaded data. See geodata_path
virtual	logical. If TRUE a virtual connection to the file is returned. This is useful if you want to extract a small area without downloading the entire raster
...	additional arguments passed to download.file

Details

var	description	unit
Al	extractable aluminum	$mg \cdot kg^{-1}$
bdr	bed rock depth	cm
clay	clay content	%
C.tot	total carbon	kg^{-1}
Ca	extractable calcium	$mg \cdot kg^{-1}$
db.od	bulk density	$kg \cdot m^3$
eCEC.f	effective cation exchange capacity	$cmol(+)kg^{-1}$
Fe	extractable iron	$mg \cdot kg^{-1}$
K	extractable potassium	$mg \cdot kg^{-1}$
Mg	extractable magnesium	$mg \cdot kg^{-1}$
N.tot	total organic nitrogen	$g \cdot kg^{-1}$
OC	Organic Carbon	$g \cdot kg^{-1}$
P	extractable phosphorus	$mg \cdot kg^{-1}$
pH.H2O	pH (H_2O)	-
sand	Sand content	%
silt	Silt content	%
S	Extractable sulfur	$mg \cdot kg^{-1}$
texture	texture class	-
wpg2	stone content	%
Zn	Extractable zinc	$mg \cdot kg^{-1}$

Value

SpatRaster

References

Tomislav Hengl, Matthew A. E. Miller, Josip Križan, Keith D. Shepherd, Andrew Sila, Milan Kilibarda, Ognjen Antonijevic, Luka Glušica, Achim Dobermann, Stephan M. Haeefe, Steve P. McGrath, Gifty E. Acquah, Jamie Collinson, Leandro Parente, Mohammadreza Sheykhmousa, Kazuki Saito, Jean-Martial Johnson, Jordan Chamberlin, Francis B.T. Silatsa, Martin Yemefack, John Wendt, Robert A. MacMillan, Ichsani Wheeler & Jonathan Crouch, 2021. African soil properties and nutrients mapped at 30 m spatial resolution using two-scale ensemble machine learning. Scientific Reports 11: 6130.

See Also

[soil_af_elements](#), [soil_af](#), [soil_world](#)

Examples

```
# downloads a large file
afph <- soil_af_isda("ph.h2o", path=tempdir(), quiet=TRUE)
```

soil_af_isda_vsi	<i>iSDA soil data for Africa, virtual connection</i>
------------------	--

Description

Virtually connect to the iSDA soil data for Africa. The spatial of these data is 30m.

For more info see:

<https://envirometrix.nl/isdasoil-open-soil-data-for-africa/>

<https://zenodo.org/search?page=1&size=20&q=iSDAsoil>

Usage

```
soil_af_isda_vsi(var)
```

Arguments

var	character. The variables name, one of: "Al", "bdr", "clay", "C.tot", "CEC", "Ca", "db.od", "eCEC.f", "Fe", "K", "Mg", "N.tot", "oc", "P", "pH.H2O", "sand", "silt", "S", "texture", "wpg2", "Zn".see Details
-----	--

Details

var	description	unit
Al	extractable aluminum	$mg \cdot kg^{-1}$
bdr	bed rock depth	cm
clay	clay content	%
C.tot	total carbon	kg^{-1}

Ca	extractable calcium	$mg \cdot kg^{-1}$
db.od	bulk density	$kg \cdot m^3$
eCEC.f	effective cation exchange capacity	$cmol(+)kg^{-1}$
Fe	extractable iron	$mg \cdot kg^{-1}$
K	extractable potassium	$mg \cdot kg^{-1}$
Mg	extractable magnesium	$mg \cdot kg^{-1}$
N.tot	total organic nitrogen	$g \cdot kg^{-1}$
OC	Organic Carbon	$g \cdot kg^{-1}$
P	Phosphorus content	$mg \cdot kg^{-1}$
pH.H2O	pH (H_2O)	-
sand	Sand content	%
silt	Silt content	%
S	Extractable sulfur	$mg \cdot kg^{-1}$
texture	texture class	-
wpg2	stone content	%
Zn	Extractable zinc	$mg \cdot kg^{-1}$

Value

SpatRaster

References

Tomislav Hengl, Matthew A. E. Miller, Josip Križan, Keith D. Shepherd, Andrew Sila, Milan Kilibarda, Ognjen Antonijevic, Luka Glušica, Achim Dobermann, Stephan M. Haefele, Steve P. McGrath, Gifty E. Acquah, Jamie Collinson, Leandro Parente, Mohammadreza Sheykhmousa, Kazuki Saito, Jean-Martial Johnson, Jordan Chamberlin, Francis B.T. Silatsa, Martin Yemefack, John Wendt, Robert A. MacMillan, Ichsani Wheeler & Jonathan Crouch, 2021. African soil properties and nutrients mapped at 30 m spatial resolution using two-scale ensemble machine learning. Scientific Reports 11: 6130.

See Also

[soil_af_elements](#), [soil_af_isda](#), [soil_world](#)

soil_af_water	<i>Soil data for water balance computation (Africa only)</i>
---------------	--

Description

Download physical soil properties data for Africa that can be used in water balance computation. The values are for a soil depth of 0 to 30 cm. The spatial resolution is 30 arc-seconds (about 1 km2), aggregated from the original 250m resolution.

For other properties see [soil_af](#), [soil_af_elements](#), [soil_af_isda](#).

For more info, see <https://www.isric.org/projects/soil-property-maps-africa-250-m-resolution>

The data have a CC-BY 4.0 NC license

Usage

```
soil_af_water(var, depth = "30cm", path, ...)
```

Arguments

var	character. Variables name such as "awcpf23" or "pwp". See Details
depth	character. Either "30cm" or "erzd" (the effective rooting zone depth of maize)
path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file

Details

var	description	unit
awcpf23	Available water capacity of the fine earth at field capacity (pF 2.3)	volumetric %
pwp	Moisture content of the fine earth at permanent wilting point (pF 4.2)	volumetric %
tetas	Moisture content of the fine earth at saturation	volumetric %
tawcpf23	Absolute total available water capacity	cm?
tawcpf23mm	Absolute total available water capacity in mm	mm
erzd	Effective root zone depth (for maize)	cm

Value

SpatRaster

See Also

[soil_af_elements](#), [soil_af_isda](#), [soil_world](#)

Examples

```
# this downloads a large file
tetaS <- soil_af_water(var="tetas", depth="erzd", path=tempdir())
```

soil_world	Global soil data
------------	------------------

Description

Download global soils data. The data are derived from the SoilGRIDS database. The data were aggregated and transformed to a longitude/latitude coordinate reference system with 30-second spatial resolution.

See <https://www.isric.org/explore/soilgrids> for more info.

data license: CC-BY 4.0

Note that currently only the "mean" values for up to 60cm depth are available.

Usage

```
soil_world(var, depth, stat="mean", name="", path, vsi=FALSE, ...)
```

Arguments

var	character. Variables name. One of: "bdod", "cfvo", "clay", "nitrogen", "ocd", "ocs", "phh2o", "sand", "silt", "soc", "wrb". See Details
depth	numeric. One of 5, 15, 30, 60, 100, 200. This is shorthand for the following depth ranges: 0-5, 5-15, 15-30, 30-60, 60-100, 100-200 cm. Ignored if var="wrb"
stat	character. One of "mean", "uncertainty", "Q0.05", "Q0.5", "Q0.95". Ignored if var="wrb"
name	character. One of "Acrisols", "Albeluvisols", "Alisols", "Andosols", "Arenosols", "Calcisols", "Cambisols", "Chernozems", "Cryosols", "Durisols", "Ferralsols", "Fluvisols", "Gleysols", "Gypsisols", "Histosols", "Kastanozems", "Leptosols", "Lixisols", "Luvisols", "Nitisols", "Phaeozems", "Planosols", "Plinthosols", "Podzols", "Regosols", "Solonchaks", "Solonetz", "Stagnosols", "Umbrisols", "Vertisols". Only used when var="wrb"
path	character. Path for storing the downloaded data. See geodata_path
vsi	logical. If TRUE, a virtual connection to the file is used in stead of a download. This allows downloading a small area only by subsequently using crop or window
...	additional arguments passed to download.file

Details

var	description	unit
bdod	Bulk density of the fine earth fraction	$kg \cdot dm^{-3}$
cec	Cation Exchange Capacity of the soil	$cmol(+)kg^{-1}$
cfvo	Vol. fraction of coarse fragments (> 2 mm)	%

nitrogen	Total nitrogen (N)	$g \cdot kg^{-1}$
phh2o	pH (H_2O)	-
sand	Sand (> 0.05 mm) in fine earth	%
silt	Silt (0.002-0.05 mm) in fine earth	%
clay	Clay (< 0.002 mm) in fine earth	%
soc	Soil organic carbon in fine earth	$g \cdot kg^{-1}$
ocd	Organic carbon density	$kg \cdot m^{-3}$
ocs	Organic carbon stocks	$kg \cdot m^{-2}$

Value

SpatRaster

References

Poggio L., de Sousa L.M., Batjes N.H., Heuvelink G.B.M., Kempen B., Ribeiro E., Rossiter D., 2021. SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty. Soil 7:217-240, 2021. doi:10.5194/soil-7-217-2021

See Also

For virtual access to the original data: [soil_world_vsi](#) For Africa: [soil_af_isda](#), [soil_af](#), [soil_af_elements](#)

Examples

```
# this downloads a large file
gph <- soil_world(var="phh2o", depth=5, path=tempdir())
```

soil_world_vsi

Global soil data, virtual connection

Description

Virtually connect to the global soilgrids data. See <https://www.isric.org/explore/soilgrids> for more info.

data license: CC-BY 4.0

Usage

```
soil_world_vsi(var, depth, stat="mean", name="")
```

Arguments

var	character. Variables name. One of: "bdod", "cfvo", "clay", "nitrogen", "ocd", "ocs", "phh2o", "sand", "silt", "soc", "wrb". See Details
depth	numeric. One of 5, 15, 30, 60, 100, 200. This is shorthand for the following depth ranges: 0-5, 5-15, 15-30, 30-60, 60-100, 100-200 cm. Ignored if var="wrb"
stat	character. One of "mean", "uncertainty", "Q0.05", "Q0.5", "Q0.95". Ignored if var="wrb"
name	character. One of 'Acrisols', 'Albeluvisols', 'Alisols', 'Andosols', 'Arenosols', 'Calcisols', 'Cambisols', 'Chernozems', 'Cryosols', 'Durisols', 'Ferralsols', 'Fluvisols', 'Gleysols', 'Gypsisols', 'Histosols', 'Kastanozems', 'Leptosols', 'Lixisols', 'Luvisols', 'Nitisols', 'Phaeozems', 'Planosols', 'Plinthosols', 'Podzols', 'Regosols', 'Solonchaks', 'Solonetz', 'Stagnosols', 'Umbrisols', 'Vertisols'. Only used when var="wrb"

Details

The below table lists the variable names, a description, and the units of the variables. Note that these units are not standard units, and are different from the data for other soil data available through this package.

var	description	unit
bdod	Bulk density of the fine earth fraction	$cg \cdot cm^{-3}$
cec	Cation Exchange Capacity of the soil	$mmol(+)kg^{-1}$
cfvo	Vol. fraction of coarse fragments (> 2 mm)	%
nitrogen	Total nitrogen (N)	$cg \cdot kg^{-1}$
phh2o	pH (H_2O)	-
sand	Sand (> 0.05 mm) in fine earth	%
silt	Silt (0.002-0.05 mm) in fine earth	%
clay	Clay (< 0.002 mm) in fine earth	%
soc	Soil organic carbon in fine earth	$dg \cdot kg^{-1}$
ocd	Organic carbon density	$hg \cdot m^{-3}$
ocs	Organic carbon stocks	$hg \cdot m^{-2}$

Value

SpatRaster

References

Poggio, L., de Sousa, L.M., Batjes, N.H., Heuvelink, G.B.M., Kempen, B., Ribeiro, E., and Rossiter, D., 2021. SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty. Soil 7:217-240, 2021. doi:10.5194/soil-7-217-2021

See Also

[soil_world](#) to download these data at 30-seconds spatial resolution.

For Africa: [soil_af_isda](#), [soil_af](#), [soil_af_elements](#)

Examples

```
ph <- soil_world_vsi(var="phh2o", depth=5)
ph
```

sp_occurrence

Download species occurrence data from GBIF

Description

Download data from the Global Biodiversity Information Facility ([GBIF](#)) data portal.

sp_genus returns a data.frame with all the species names associated with a genus.

sp_occurrence downloads occurrence records for a single species.

You can check the number of records returned by using the option download=FALSE.

Note that the you can only download up to record number 100,000. To avoid getting more than 100,000 records, you can do separate queries for different geographic areas. This has been automated in sp_occurrence_split. This function recursively splits the area of interest into smaller areas until the number of records in an area is less than 50,000. It then downloads these records and saves them in a folder called "gbif". After all areas have been evaluated, the data are combined into a single file and returned as a data.frame or SpatVector). If the function is interrupted, it can be run again, and it will resume where it left off.

If you want to download data for an entire genus, first run sp_genus and then download data for the returned species names one by one.

Before using this function, please first check the GBIF [data use agreement](#) and see the note below about how to cite these data.

Usage

```
sp_genus(genus, simple=TRUE, ...)
```

```
sp_occurrence(genus, species="", ext=NULL, args=NULL, geo=TRUE,
  removeZeros=FALSE, download=TRUE, ntries=5, nrecs=300,
  start=1, end=Inf, fixnames=TRUE, sv=FALSE, ...)
```

```
sp_occurrence_split(genus, species="", path=".", ext=c(-180,180,-90,90),
  args=NULL, geo=TRUE, removeZeros=FALSE, ntries=5, nrecs=300,
  fixnames=TRUE, prefix=NULL, sv=FALSE, ...)
```

Arguments

genus	character. genus name
species	character. species name
ext	SpatExtent object to limit the geographic extent of the records. A SpatExtent can be created using functions like ext and draw

args	character. Additional arguments to refine the query. See query parameters in http://gbif.org/developer/occurrence for more details
geo	logical. If TRUE, only records that have a georeference (longitude and latitude values) will be downloaded
removeZeros	logical. If TRUE, all records that have a latitude OR longitude of zero will be removed if geo==TRUE, or set to NA if geo==FALSE. If FALSE, only records that have a latitude AND longitude that are zero will be removed or set to NA
download	logical. If TRUE, records will be downloaded, else only the number of records will be shown
ntries	integer. How many times should the function attempt to download the data, if an invalid response is returned (perhaps because the GBIF server is very busy)
nrecs	integer. How many records to download in a single request (max is 300)?
start	integer. Record number from which to start requesting data
end	integer. Last record to request
fixnames	If TRUE a few unwieldy and poorly chosen variable names are changed as follows. "decimalLatitude" to "lat", "decimalLongitude" to "lon", "stateProvince" to "adm1", "county" to "adm2", "countryCode" to "ISO2". The names in "country" are replaced with the common (short form) country name, the original values are stored as "fullCountry"
path	character. Where should the data be downloaded to (they will be put in a subdirectory "gbif")?
prefix	character. prefix of the downloaded filenames (best left NULL, the function will then use "genus_species")
simple	logical. If TRUE, a vector the accepted species names are returned. Otherwise a data.frame with much more information is returned
sv	logical. If TRUE, a SpatVector rather than a data frame is returned.
...	additional arguments passed to download.file

Value

data.frame, or SpatVector if sv=TRUE

Note

Under the terms of the GBIF data user agreement, users who download data agree to cite a DOI. Citation rewards data-publishing institutions and individuals and provides support for sharing open data [1][2]. You can get a DOI for the data you downloaded by creating a **"derived" dataset**. For this to work, you need to keep the "datasetKey" variable in your dataset.

References

<https://gbif.org/occurrence> <https://gbif.org/derived-dataset/about>

Examples

```
sp_occurrence("solanum", "acaule", download=FALSE)
g <- sp_occurrence("Solanum", "acaule", geo=TRUE, end=5)

sp_occurrence("Batrachoseps", "luciae", down=FALSE)
g <- sp_occurrence("Batrachoseps", "luciae", geo=TRUE, end=5)

plot(g[, c("lon", "lat")])

## args
a1 <- sp_occurrence("Elgaria", "multicarinata",
  args="recordNumber=Robert J. Hijmans RH-2")
a2 <- sp_occurrence("Batrachoseps", "luciae",
  args=c("year=2023", "identifiedBy=Anthony Ye"))

## year supports "range queries"
a3 <- sp_occurrence("Batrachoseps", "luciae",
  args=c("year=2020,2023", "identifiedBy=Kuoni W"))
table(a3[,c("year")])
```

travel_time	<i>Travel time to a city or port</i>
-------------	--------------------------------------

Description

Download global travel time to a city or port data on rasters at a 30 arc-seconds (about 1 km2) resolution.

Usage

```
travel_time(to="city", size=1, up=FALSE, path, ...)
```

Arguments

to	character. "city" or "port"
size	positive integer indicating the size of the city or port. Can be between 1 and 9 if to="city" or between 1 and 5 if to="port". See Details
up	logical. If TRUE the travel time to a city of the size chosen or larger is returned
path	character. Path for storing the downloaded data. See geodata_path
...	additional arguments passed to download.file

Details

Description of the the size argument.

to="city"

size	Inhabitants
1	5,000,000 to 50,000,000
2	1,000,000 to 5,000,000
3	500,000 to 1,000,000
4	200,000 to 500,000
5	100,000 to 200,000
6	50,000 to 100,000
7	20,000 to 50,000
8	10,000 to 20,000
9	5,000 to 10,000

to="port"

size	Description	Number of ports
1	Large	160
2	Medium	361
3	Small	990
4	Very small	2,153
5	Any	3,778

Value

SpatRaster

References

Nelson, A., D.J. Weiss, J. van Etten, A. Cattaneo, T.S. McMenomy & J. Koo, 2019. A suite of global accessibility indicators. Scientific Data 6: 266. doi:10.1038/s41597-019-0265-5

Version 3 (2019-05-15) from https://figshare.com/articles/dataset/Travel_time_to_cities_and_ports_in_the_year_2015/7638134/3

world

Administrative boundaries

Description

Get the borders for all the countries in the world. Data are read from files that are downloaded if necessary.

Usage

```
world(resolution=5, level=0, path, version="latest", ...)
```

Arguments

resolution	integer between 1 and 5 indicating the level of detail. 1 is high 5 is low
level	numeric. The level of administrative subdivision requested. (starting with 0 for country, then 1 for the first level of subdivision). Only level 0 is currently available
path	character. Path for storing the downloaded data. See geodata_path
version	character. Only "3.6" is currently supported
...	additional arguments passed to download.file

Details

The data are from <https://gadm.org>

Value

SpatVector

See Also

[gadm](#)

Examples

```
w <- world(path=tempdir())
```

worldclim

WorldClim climate data

Description

Download climate data from WorldClim version 2.1. See Details for variables and units.

Usage

```
worldclim_global(var, res, path, version="2.1", ...)
worldclim_country(country, var, path, version="2.1", ...)
worldclim_tile(var, lon, lat, path, version="2.1", ...)
```

Arguments

var	character. Valid variables names are "tmin", "tmax", "tavg", "prec", "wind", "vapr", and "bio"
res	numeric. Valid resolutions are 10, 5, 2.5, and 0.5 (minutes of a degree)
path	character. Path for storing the downloaded data. See geodata_path
country	character. Country name or code
lon	numeric. Longitude
lat	numeric. Latitude
version	character or numeric. WorldClim version number. Only "2.1" supported at the moment
...	additional arguments passed to download.file

Details

These are the WorldClim monthly average climate data.

Variable	Description	Unit
tmin	minimum temperature	°C
tmax	maximum temperature	°C
tavg	average temperature	°C
prec	total precipitation	mm
srad	incident solar radiation	$\text{kJ} \cdot \text{m}^{-2} \cdot \text{day}^{-1}$
wind	wind speed (2 m above the ground)	$\text{m} \cdot \text{s}^{-1}$
vapr	vapor pressure	kPa

Value

SpatRaster

See Also

<https://www.worldclim.org/>

Examples

```
lux <- worldclim_country("Luxembourg", var="tmin", path=tempdir())
```

Index

* **spatial**

sp_occurrence, [27](#)

bio_oracle, [2, 3](#)

cmip6, [4](#)

cmip6_tile, [2](#)

cmip6_tile (cmip6), [4](#)

cmip6_world, [2](#)

cmip6_world (cmip6), [4](#)

country_codes, [2, 5](#)

crop, [24](#)

crop_calendar_sacks, [2, 6](#)

crop_monfreda, [2, 7](#)

crop_spam, [2, 8](#)

cropland, [2, 9](#)

download.file, [3, 5–12, 14–18, 20, 23, 24, 28, 29, 31, 32](#)

draw, [27](#)

elevation, [10](#)

elevation_30s, [2](#)

elevation_30s (elevation), [10](#)

elevation_3s, [2](#)

elevation_3s (elevation), [10](#)

elevation_global, [2](#)

elevation_global (elevation), [10](#)

ext, [27](#)

footprint, [3, 11](#)

gadm, [2, 12, 31](#)

geodata (geodata-package), [2](#)

geodata-package, [2](#)

geodata_path, [3, 5–12, 13, 14–18, 20, 23, 24, 29, 31, 32](#)

landcover, [3, 9, 11, 13, 14](#)

monfredaCrops (crop_monfreda), [7](#)

osm, [3, 14](#)

population, [3, 15](#)

rice_calendar, [16](#)

sacksCrops (crop_calendar_sacks), [6](#)

soil_af, [3, 16, 19, 21, 22, 25, 26](#)

soil_af_elements, [3, 18, 18, 21–23, 25, 26](#)

soil_af_isda, [3, 16, 18, 19, 19, 22, 23, 25, 26](#)

soil_af_isda_vsi, [21](#)

soil_af_water, [3, 22](#)

soil_world, [3, 16, 19, 21–23, 24, 26](#)

soil_world_vsi, [3, 18, 25, 25](#)

sp_genus (sp_occurrence), [27](#)

sp_occurrence, [3, 27](#)

sp_occurrence_split (sp_occurrence), [27](#)

spamCrops (crop_spam), [8](#)

travel_time, [3, 29](#)

vrt, [5](#)

window, [24](#)

world, [3, 12, 30](#)

worldclim, [31](#)

worldclim_country, [3](#)

worldclim_country (worldclim), [31](#)

worldclim_global, [3](#)

worldclim_global (worldclim), [31](#)

worldclim_tile, [3](#)

worldclim_tile (worldclim), [31](#)